

# Long term data records of stratospheric composition from the GOZCARDS Project: Variations in HCl, O<sub>3</sub>, and H<sub>2</sub>O




L. Froidevaux<sup>1</sup>, J. Anderson<sup>2</sup>, H.-J. (Ray) Wang<sup>3</sup>, R. Fuller<sup>1</sup>, N. J. Livesey<sup>1</sup>,  
H. C. Pumphrey<sup>4</sup>, S. Davis<sup>5</sup>, K. Rosenlof<sup>5</sup>, R. McPeters<sup>6</sup>, S. M. Frith<sup>7</sup>,  
J. Wild<sup>8</sup>, J. M. Russell<sup>2</sup>, P. F. Bernath<sup>9</sup>, K. A. Walker<sup>10</sup>,  
J. M. Zawodny<sup>11</sup>, and L. W. Thomason<sup>11</sup>

1. Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA
2. Hampton University, Hampton, VA, USA
3. Georgia Institute of Technology, Atlanta, GA, USA
4. University of Edinburgh, Edinburgh, Scotland
5. NOAA ESRL Chemical Sciences Division, Boulder, CO, USA
6. NASA Goddard Space Flight Center, Greenbelt, MD, USA
7. SSAI; NASA Goddard Space Flight Center, MD, USA
8. Wyle ST&E, NOAA/NCEP Climate Prediction Center, MD USA
9. Old Dominion University, Norfolk, VA, USA
10. University of Toronto, Toronto, Canada
11. NASA Langley Research Center, Hampton, VA, USA

**GOZCARDS: Global OZone Chemistry And Related trace gas**  
**Data records for the Stratosphere**  
*part of the NASA MEaSUREs program*

# Satellite/Instrument Timeline and data versions



**Timeline of satellite missions and instruments considered for the GOZCARDS project and the creation of a stratospheric composition Earth System Data Record (ESDR).**

## Common Grids

- Mixing ratios (time, lat., p)
  - Monthly zonal averages
  - 10 degree latitude bins
  - $p(i) = 1000/10^{(-i/6)}$   $i=0, 1, 2, \dots$   
(same as UARS pressure grid)

## Data Versions (for creating merged data records)


	O <sub>3</sub>	H <sub>2</sub> O	HCl
SAGE I	5.9	-	-
SAGE II	6.2	-	-
UARS MLS	5	6	-
HALOE	19	19	19
ACE-FTS	2.2u	2.2	2.2
Aura MLS	2.2	3.3	3.3

## netCDF source files & merged files

- > include mean values, but also
  - std. deviations, std. errors,
  - + info on local time, SZA,
  - days used each month
  - + offsets applied to each source dataset

# GOZCARDS methodology for merging datasets (H<sub>2</sub>O, HCl)


- Merging process uses a bias correction method to tie time series together into one longer-term series
  - > obtain average offsets during overlap time period(s)
  - > make use of the good temporal coverage provided by MLS, and iterate using ACE-FTS and HALOE consecutively (weighting = 1/3 for each)
  - > result (in H<sub>2</sub>O example below) is equivalent to using 3-way average during overlap period



**Methodology for HCl and H<sub>2</sub>O is basically identical**


(but for H<sub>2</sub>O, add UARS MLS as an extra step; also, ignore Aura MLS HCl for  $p < 10$  hPa.)

# Examples: GOZCARDS Offsets for H<sub>2</sub>O




# GOZCARDS methodology for O<sub>3</sub>

- For each individual satellite dataset, first calculate **monthly zonal means** (in ppmv) **for each 10° latitude bin and pressure level** (~2.5 km spacing) with careful screenings.
- Adjust datasets to a **reference level** that is equal to or based on **SAGE-II average**  
> then, average the adjusted (and collocated) datasets to derive a merged ozone record.



- Note:** above 3.2 hPa, use **adjusted HALOE (HALOE\*)** instead of SAGE-II as reference, due to anomalous NCEP temperature trends after June 2000 (see next page)




(Left). O<sub>3</sub> from **SAGE-I/II**, **HALOE**, **UMLS/AMLS**, and **ACE-FTS** between 0 and 10°S at 46.4 hPa.  
(Right). Adjusted source datasets and merged time series for O<sub>3</sub> in same lat./p bin as left panel.

Monthly zonal mean ozone from **SAGE**, **HALOE**, **MLS**, and **ACE**

- Issue (mainly for upper strat. after mid-2000) [McLinden et al., 2011]

T-related (NCEP) drifts impact ozone time series for SAGE II data converted to VMR/p grid




# Comparisons between GOZCARDS and SWOOSH

- The **Stratospheric Water and OzOne Satellite Homogenized (SWOOSH)** database (*from preliminary version - Sean Davis, Karen Rosenlof, NOAA*)
  - **Datasets used**
    - SAGE-II, UARS MLS, HALOE, Aura-MLS
      - > notes: SAGE II H<sub>2</sub>O not used in GOZCARDS (channel drift issue concerns)
      - UARS MLS not (yet) used in SWOOSH O<sub>3</sub> data
  - **Report monthly zonal means**  
(both latitude/pressure and equivalent latitude/PV surfaces)
    - Volume mixing ratios (monthly means)
      - 18 latitude bins (every 10°) [also report data in 2.5°bins]
      - Aura MLS (v3.3) pressure levels
  - **Merging method for SWOOSH**
    - **Use Aura-MLS as reference**
      - > **GOZCARDS uses SAGE II for O<sub>3</sub> and avg [HALOE, AMLS, ACE-FTS] for H<sub>2</sub>O**
    - Calculate offsets based on collocated profile pairs (within latitude bins)
- Differences above (+ other diffs.) in source datasets and merging methods  
→ we do not expect a “perfect match” for GOZCARDS versus SWOOSH
  - but this is a **useful cross-check for O<sub>3</sub> and H<sub>2</sub>O results**  
(for both the GOZCARDS and SWOOSH teams)




# O<sub>3</sub>: Comparisons between GOZCARDS and SWOOSH (V2.0) (1984-2010)

**SWOOSH and GOZCARDS O<sub>3</sub> series at 10 hPa for 40S-50S**




relative trend in differences  
0.02 ppmv/decade (~ 0.3%/decade)

**SWOOSH and GOZCARDS O<sub>3</sub> series at 3.1 hPa for 20S-30S**



relative trend in differences:  
0.15 ppmv/decade (~2%/decade)

Mean diffs. are  
< 5% in most of  
stratosphere  
- larger % diff. in tropical  
UTLS (where SAGE II  
& Aura MLS differ more)




Trend differences in  
upper strat. arise  
from

- different source data  
& references
- different treatment  
of SAGE II after  
mid-2000.









# Average differences of zonal means versus averages of coincidences SAGE II versus Aura MLS Ozone (2004-2005)



- Average offsets are not very dependent on the method used, although some larger differences can exist in more localized latitude bins.
- Diurnal effect plays a role in upper stratosphere / lower mesosphere
  - > nighttime Aura MLS values are used above for the coincident method
  - gives better average agreement with SAGE II twilight data

# O<sub>3</sub> Comparisons: SWOOSH versus GOZCARDS anomalies (1984 to 2010)

## Correlation Coefficient



## Trend of differences:

< ~1%/dec,  $p > 3$  hPa

2-3%/dec,  $p < 3$  hPa


- also larger diffs.  
in UTLS region

Grey shading  
indicates where  
trends are not  
significantly  
different from zero  
(3-sigma test)

# O<sub>3</sub> Comparisons: homogenized SBUV versus GOZCARDS anomalies (1984 to 2010)

**NASA Profile MOD** (from R. McPeters, S. Frith, et al.)


- No offset corrections between different NOAA satellites



Shading → trends not significantly different from zero (3-sigma test).


**NOAA-MA-SBUV** (from J. Wild et al.)

- Offset corrections are applied between different NOAA satellites





## Deseasonalized Anomalies and diffs. (SWOOSH - GOZCARDS)

Corr. Coef.




**Slopes (ppmv/decade)**



# H<sub>2</sub>O: Comparisons between GOZCARDS and SWOOSH

## H<sub>2</sub>O Anomalies (tracking the variability)




Yellow shading indicates slopes that are not statistically significant at the 3 sigma level. Contours are in 1 %/decade increments.


# Temporal variations: A few more examples

(slight amount of interpolation/smoothing applied)

## HCl



HCl Column ( $\times 10^{15}$  mol/cm<sup>2</sup>)



Global merged HCl at 1 hPa ( $\approx$  total chlorine)


Witnessing (measuring) the rise and fall of the evil (chlorine) empire...

Column HCl (100 to 1 hPa)

- GOZCARDS trends appear consistent with ground-based total column results ( $\sim -0.8\%/yr$ )  
(Jungfraujoch column data shown above)



# Temporal variations: A few more examples




## H<sub>2</sub>O: interesting LS variations


- as observed/discussed in literature (and at this meeting)
- but will the post-2004 rise continue?
- > implications for T, circulation, and climate

## O<sub>3</sub>: LS ozone recovery?

- requires detailed analyses




## O<sub>3</sub>: Is upper stratospheric ozone on its way back up?





# Temporal variations: A few more examples




## H<sub>2</sub>O: interesting LS variations


- as observed/discussed in literature (and at this meeting)
- but will the post-2004 rise continue?
- > implications for T, circulation, and climate

## O<sub>3</sub>: LS ozone recovery?

- requires detailed analyses



O<sub>3</sub>: Is upper stratospheric ozone on its way back up?



# GOZCARDS Status, Upcoming Work


- Deliver HCl, O<sub>3</sub>, H<sub>2</sub>O, & T (MERRA) to GES DISC for access (starting this month)
- Write overview paper + GES DISC README guide (latter is essentially done)
- Continue GOZCARDS work on other species

HF, ClO, HNO<sub>3</sub>, CH<sub>4</sub>, N<sub>2</sub>O, NO<sub>2</sub>, NO, ClO<sub>x</sub>, NO<sub>x</sub>

- > some involve fewer instruments (may seem simpler, but also get less overlapping data)
- > each species poses its own challenges

**A lesson for the future:** especially for occultation data (e.g., SAGE or ACE follow-on), good to have > 2 years of overlap with ongoing missions (if possible...)

Merged HCl Data at 46 hPa



- We expect community feedback, once GOZCARDS goes public  
→ some iteration possible
- GOZCARDS data records are generally as close as possible to the original data (after screening, despiking,...)  
→ **there are** (sometimes large) **data gaps**
- Users may want to smooth or fit data in different ways → trends, etc... (e.g., for SI<sup>2</sup>N)  
> 'smart' sampling of models is useful
- Short-term portions of the series are only as good as the input datasets, but a long-term carefully produced ESDR should empower the user community to pursue further research.